

COMPLIMENTARY RECEPTION

Finding the Zen in Managing Atopic Dermatitis

February 26, 2026
Thursday

Reception: 6:00 PM
Event: 7:00 PM

Embassy Suites Murfreesboro
1200 Conference Center Blvd.

Join us for an engaging program that takes a real-world look at Janus kinase (JAK) inhibitors and how they fit into everyday management of itchy dogs. We'll break down how these drugs work, when to use them, and which cases benefit most. You'll also get a clear, clinician-friendly overview of today's treatment options for canine atopic dermatitis and how to choose the right approach for each patient. **Attendees will receive one hour of CE.**



Joya Griffin, DVM, DACVD

Dr. Griffin is a board-certified veterinary dermatologist practicing with Animal Dermatology Clinic in Louisville. A Cornell graduate and ACVD Diplomate, she is an internationally recognized speaker and the star of Pop Goes the Vet with Dr. Joya on NatGeoWILD and Disney+.

Register



IMPORTANT SAFETY INFORMATION

INDICATIONS

Zenrelia is indicated for control of pruritus associated with allergic dermatitis and control of atopic dermatitis in dogs at least 12 months of age.

Read the entire package insert, including the Boxed Warning before using this drug.

For full prescribing information call 1-888-545-5973 or visit <http://www.elancolabels.com/us/zenrelia>

WARNING: INADEQUATE IMMUNE RESPONSE TO VACCINES. Based on results of the vaccine response study, dogs receiving Zenrelia are at risk of an inadequate immune response to vaccines. Discontinue Zenrelia for at least 28 days to 3 months prior to vaccination and withhold Zenrelia for at least 28 days after vaccination. Dogs should be up to date on vaccinations prior to starting Zenrelia. Do not use in dogs less than 12 months old or dogs with a serious infection. Monitor dogs for infections because Zenrelia may increase susceptibility to infections, including adenoviral hepatitis and pancreatitis. Neoplastic conditions (benign and malignant) were observed during clinical studies. Consider the risks and benefits of treatment in dogs with a history of recurrence of these conditions. The most common adverse reactions were vomiting, diarrhea and lethargy. Zenrelia has not been evaluated in breeding, pregnant, or lactating dogs and concurrent use with glucocorticoids, cyclosporine, or other systemic immunosuppressive agents has not been tested. For full prescribing information see [package insert](#).